

Shiro Miyayama, M.D.

Nationality: Japanese

Affiliation: Department of Diagnostic Radiology, Fukuiken Saiseikai Hospital

Address: 7-1, Funabashi, Wadanaka-cho, Fukui 918-8503, Japan

Education:

1980-1986 Tottori University School of Medicine

Working Experience:

1986-1988 Department of Radiology, Kanazawa University School of Medicine

(Resident)

1988-1989 Department of Diagnostic Radiology, Fukuiken Saiseikai Hospital

1989-1990 Department of Radiology, Toyama Red Cross Hospital

1990-1991 Department of Radiology, Kaga Central Hospital

1991-1995 Department of Radiology, Kanazawa University School of Medicine (Staff)

1995- Department of Diagnostic Radiology, Fukuiken Saiseikai Hospital

(Department head)

Specialty: Interventional Radiology, Abdominal imaging

Membership of Academic Society: JSIR, JRC, SIR, CIRSE, etc.

Specialty Board Certification: Diagnostic Radiologist (JRS), Interventional Radiologist (JSIR)

Positions in Academic Society: JSIR (Secretary), JRC (Delegate), The Japanese Society for Transcatheter Hepatic Arterial Embolization (Vice Secretary), The Japanese Society of Endoluminal Metallic Stents & Grafts (Secretary), etc.

Award: JSIR best paper award (2001), JVIR distinguish reviewer award (2016), JVIR top reviewer award (2017), etc.

Articles published in the major English journals (First author only)

1. Miyayama S, Matsui O, Kadoya M, Yoshikawa J, Gabata T, Kitagawa K, Arai K, Takashima T. Malignant esophageal stricture and fistula: palliative treatment with

- polyurethane-covered Gianturco stent. *J Vasc Interv Radiol* 1995; 6 (2):242-248.
2. Miyayama S, Matsui O, Kamimura R, Kakuta K, Takashima T. Partially covered Gianturco stent for trachemobronchial stricture caused by intraluminal tumor. *Cardiovasc Intervent Radiol* 1997; 20 (1):60-62.
 3. Miyayama S, Matsui O, Terayama N, Tatsu H, Yamamoto T, Takashima T. Covered Gianturco stents for malignant biliary obstruction: preliminary clinical evaluation. *J Vasc Interv Radiol* 1997; 8 (4):641-648.
 4. Miyayama S, Matsui O, Ueda K, Kifune K, Yamashiro M, Yamamoto T, Komatsu T, Kumano T. Hemodynamics of small hepatic focal nodular hyperplasia: evaluation with single-level dynamic CT during hepatic arteriography. *AJR Am J Roentgenol* 2000; 174 (6):1567-159.
 5. Miyayama S, Matsui O, Kifune K, Yamashiro M, Yamamoto T, Kitagawa K, Kasahara Y, Asada Y, Iida Y, Miura S. Malignant colonic obstruction due to extrinsic tumor: palliative treatment with a self-expanding nitinol stent. *AJR Am J Roentgenol* 2000; 175 (6):1631-1637.
 6. Miyayama S, Matsui O, Akakura Y, Yamamoto T, Fujinaga Y, Koda W, Kawai K, Notsumata K, Toya D, Tanaka N. Use of a catheter with a large side hole for selective catheterization of the inferior phrenic artery. *J Vasc Interv Radiol* 2001; 12 (4):497-499.
 7. Miyayama S, Matsui O, Akakura Y, Yamamoto T, Nishida H, Yoneda K, Kawai K, Nishijima H. Hepatocellular carcinoma with blood supply from omental branches: treatment with transcatheter arterial embolization. *J Vasc Interv Radiol* 2001; 12 (11):1285-1290.
 8. Miyayama S, Matsui O, Akakura Y, Yamamoto T, Nishida H, Minami T, Kozaka K, Tawaraya K, Kasahara Y. Percutaneous cholecystochoedocostomy for cholecystitis and cystic duct obstruction in gallbladder carcinoma. *J Vasc Interv Radiol* 2003; 14 (2):261-264.
 9. Miyayama S, Matsui M, Nishida H, Yamamori S, Minami T, Shinmura R, Kozaka K, Notsumata K, Toya D, Tanaka N, Mitsui T, Nishijima H. Transcatheter arterial chemoembolization for unresectable hepatocellular carcinoma fed by the cystic artery. *J Vasc Interv Radiol* 2003; 14 (9):1155-1161.
 10. Miyayama S, Matsui M, Akakura Y, Yamamoto T, Nishida N, Yoneda K, Kawai K, Murakami S. Abdominal aortic aneurysm with a dilated common iliac artery: treatment using a handmade bifurcated stent-graft with a wide iliac limb end. *Cardiovasc Intervent Radiol* 2003; 26 (6):511-515.
 11. Miyayama S, Matsui O, Yamamoto T, Akakura Y, Yamamori S, Minami T, Kozaka K,

- Mitsui T. Intrahepatic accessory spleen: evaluation by CT arteriography. *Abdom Imaging* 2003; 28 (6):862-865.
12. Miyayama S, Nishida H, Matsui O. Peritumoral fatty infiltration of the liver associated with venous drainage from metastatic liver tumor. *AJR Am J Roentgenol* 2004; 182 (2):533-534.
 13. Miyayama S, Matsui O, Akakura Y, Yamamoto T, Nishida H, Yoneda K, Kawai K, Toya D, Tanaka N, Mitsui T, Asada Y. Efficacy of covered metallic stents in the treatment of unresectable malignant biliary obstruction. *Cardiovasc Intervent Radiol* 2004; 27 (4): 349-354.
 14. Miyayama S, Matsui O, Taki K, Minami T, Ito C, Shinmura R, Takamatsu S, Kobayashi M, Notsumata K, Toya D, Tanaka N. Transcatheter arterial chemoembolization for hepatocellular carcinoma fed by the reconstructed inferior phrenic artery: anatomical and technical analysis. *J Vasc Interv Radiol* 2004; 15 (8): 815-823.
 15. Miyayama S, Matsui O, Taki K, Minami T, Ito C, Shinmura R, Takamatsu S, Kobayashi M, Notsumata K. Combined use of an occlusion balloon catheter and a microcatheter for embolization of the unselectable right inferior phrenic artery supplying hepatocellular carcinoma. *Cardiovasc Intervent Radiol* 2004; 27 (6): 677-681.
 16. Miyayama S, Matsui O, Taki K, Minami T, Ryu Y, Ito C, Nakamura K, Inoue D, Takamatsu S. Arterial blood supply to the posterior aspect of segment IV of the liver from the caudate branch: demonstration at CT after iodized oil injection. *Radiology* 2005; 237 (3):1110-1114.
 17. Miyayama S, Matsui O, Taki K, Minami T, Shinmura R, Ito C, Takamatsu S, Kobayashi M, Ushioji Y. Occluded Brescia-Cimino hemodialysis fistulas: endovascular treatment with both brachial arterial and venous access using the pull-through technique. *Cardiovasc Intervent Radiol* 2005; 28 (6):806-812.
 18. Miyayama S, Matsui O, Taki K, Minami T, Ryu Y, Ito C, Nakamura K, Inoue D, Notsumata K, Toya D, Tanaka N, Mitsui T. Extrahepatic blood supply to hepatocellular carcinoma: angiographic demonstration and transcatheter arterial chemoembolization. *Cardiovasc Intervent Radiol* 2006; 29 (1):39-48.
 19. Miyayama S, Matsui O, Zen Y, Yamashiro M, Ryu Y, Minami T, Notsumata K, Tanaka N. Focal hepatic lesions mimicking cavernous hemangioma supplied by the portal vein. *Hepato Res* 2006; 36 (1):70-73.
 20. Miyayama S, Matsui O, Taki K, Minami T, Ito C, Shinmura R, Takamatsu S, Kobayashi M, Toya D, Mitsui T. Bile duct disruption after blunt hepatic trauma:

- treatment with percutaneous repair. *J Trauma* 2006; 60 (3):640-643.
21. Miyayama S, Matsui O, Yamashiro M, Ryu Y, Kaito K, Ozaki K, Takeda T, Yoneda Y, Notsumata K, Toya D, Tanaka N, Mistui T. Ultraslective transcatheter arterial chemoembolization with a 2-F tip microcatheter for small hepatocellular carcinomas: relationship between local tumor recurrence and visualization of portal vein with iodized oil. *J Vasc Interv Radiol* 2007; 18 (3):365-376.
 22. Miyayama S, Matsui O, Yamashiro M, Ryu Y, Takata H, Takeda T, Aburano H, Shigenari N. Visualization of hepatic lymphatic vessels during transcatheter arterial chemoembolization for hepatocellular carcinoma. *J Vasc Interv Radiol* 2007; 18 (9):1111-1117.
 23. Miyayama S, Matsui O, Yamashiro M, Ryu Y, Takata H, Takeda T, Aburano H, Shigenari N. Iodized oil accumulation in the hypovascular tumor portion of early-stage hepatocellular carcinoma after ultraslective transcatheter arterial chemoembolization. *Hepatol Int* 2007; 1 (4):451-459.
 24. Miyayama S, Yamashiro M, Okuda M, Takeda T, Aburano H, Shigenari N, Aotake Y, Matsui O. Balloon angioplasty for traumatic popliteal artery disruption: sustained patency at 9 years. *J Endovasc Ther* 2008; 15 (1):133-134.
 25. Miyayama S, Yamashiro M, Takeda T, Aburano H, Komatsu T, Sanada T, Kosaka S, Toya D, Matsui O. Acute cholecystitis caused by malignant cystic duct obstruction: treatment with metallic stent placement. *Cardiovasc Intervent Radiol* 2008; 31 (Suppl 2):S221-226.
 26. Miyayama S, Yamashiro M, Okuda M, Aburano H, Shigenari N, Morinaga K, Matsui O. Creation of a cleft in an angiography catheter to facilitate catheterization of branches of the aorta arising at an acute angle. *J Vasc Interv Radiol* 2008; 19 (12):1769-1771.
 27. Miyayama S, Yamashiro M, Okuda M, Yoshie Y, Shigenari N, Igarashi S, Nakashima Y, Matsui O. Usefulness of cone-beam computed tomography during ultraslective chemoembolization for small hepatocellular carcinomas that cannot be demonstrated on angiography. *Cardiovasc Intervent Radiol* 2009; 32 (2):255-264.
 28. Miyayama S, Mitsui O, Zen Y, Sudo Y, Yamashiro M, Okuda M, Yoshie Y, Sanada T, Notsumata K, Tanaka N, Matsui O. Histopathologic findings after ultraslective transcatheter arterial chemoembolization for hepatocellular carcinoma. *Hepatol Res* 2009; 39 (4):374-381.
 29. Miyayama S, Yamashiro M, Okuda M, Aburano H, Shigenari N, Morinaga K, Matsui O. Anastomosis between the hepatic artery and the extrahepatic collateral

- of between extrahepatic collaterals: observation on angiography. *J Med Imag Radiat Oncol* 2009; 53 (3):271-282.
30. Miyayama S, Matsui O, Yamashiro O, Ryu Y, Takata H, Takeda T, Aburano H, Shigenari N. Detection of hepatocellular carcinoma by CT during arterial portography using a cone-beam CT technology: comparison with conventional CTAP. *Abdom Imaging* 2009; 34 (4):502-506.
 31. Miyayama S, Yamashiro M, Okuda M, Yoshie Y, Shigenari N, Igarashi S, Nakashima Y, Matsui O. Hepatocellular carcinoma supplied by the right lumbar artery. *Cardiovasc Intervent Radiol* 2010; 33 (1):53-60.
 32. Miyayama S, Yamashiro M, Okuda M, Yoshie Y, Nakashima Y, Ikeno H, Orito N, Matsui O. The march of extrahepatic collaterals: analysis of blood supply to hepatocellular carcinoma located in the bare area of the liver after chemoembolization. *Cardiovasc Intervent Radiol* 2010; 33 (3):513-522.
 33. Miyayama S, Yamashiro M, Okuda M, Yoshie Y, Nakashima Y, Ikeno H, Orito N, Notsumata K, Watanabe H, Toya D, Tanaka N, Matsui O. Main bile duct stricture occurring after transcatheter arterial chemoembolization for hepatocellular carcinoma. *Cardiovasc Intervent Radiol* 2010; 33 (6):1168-1179.
 34. Miyayama S, Yamashiro M, Okuda M, Yoshie Y, Nakashima Y, Ikeno H, Orito N, Matsui O. Detection of corona enhancement of hypervascular hepatocellular carcinoma by C-arm dual-phase cone-beam CT during hepatic arteriography. *Cardiovasc Intervent Radiol* 2011; 34 (1):81-86.
 35. Miyayama S, Yamashiro M, Yoshie Y, Okuda M, Nakashima Y, Ikeno H, Orito N, Matsui O. Inferior phrenic arteries: angiographic anatomy, variations, and catheterization technique for transcatheter arterial chemoembolization. *Jpn J Radiol* 2010; 28 (7):502-511.
 36. Miyayama S, Yamashiro M, Yoshie Y, Okuda M, Nakashima Y, Ikeno H, Orito N, Ueda N, Kato T, Ushioji Y, Matsui O. Technical success rates and long-term patency of endovascular treatment for occluded native hemodialysis fistulas: comparison between thrombotic occlusion and nonthrombotic occlusion. *Jpn J Radiol* 2010; 28 (7):512-519.
 37. Miyayama S, Yamashiro M, Okuda M, Yoshie Y, Sugimori N, Igarashi S, Nakashima Y, Notsumata K, Toya D, Tanaka N, Matsui O. Chemoembolization for the treatment of large hepatocellular carcinoma. *J Vasc Interv Radiol* 2010; 21 (8):1226-1234.
 38. Miyayama S, Yamashiro M, Yoshie Y, Nakashima Y, Ikeno H, Orito N, Yoshida M, Matsui O. Hepatocellular carcinoma in the caudate lobe of the liver: variations of

- its feeding branches on arteriography. *Jpn J Radiol* 2010; 28 (8):555-562.
39. Miyayama S, Yamashiro Y, Hattori Y, Orito N, Matsui K, Tsuiji K, Yoshida M, Matsui O. Microcoil embolization during abdominal vascular interventions through microcatheters with a tip of 2 French or less. *Jpn J Radiol* 2011; 29 (4):286-290.
 40. Miyayama S, Yamashiro M, Hattori Y, Orito N, Matsui K, Tsuiji K, Yoshida M, Matsui O. Efficacy of cone-beam computed tomography during transcatheter arterial chemoembolization for hepatocellular carcinoma. *Jpn J Radiol* 2011; 29 (6):371-377.
 41. Miyayama S, Yamashiro M, Hattori Y, Orito N, Matsui K, Tsuiji K, Yoshida M, Yoshida M, Kikuchi Y, Tanaka T, Tsuda G, Matsui O. Usefulness of C-arm CT during superselective infusion chemotherapy for advanced head and neck carcinoma. *J Med Imag Radiat Oncol* 2011; 55 (4):368-372.
 42. Miyayama S, Matsui O, Zen Y, Masashi Y, Hattori Y, Orito N, Matsui K, Tsuiji K, Yoshida M, Sudo Y. Portal blood supply to locally progressed hepatocellular carcinoma after transcatheter arterial chemoembolization: observation on CT during arterial portography. *Hepatol Res* 2011; 41 (9):853-866.
 43. Miyayama S, Yamashiro M, Hattori Y, Orito N, Matsui K, Tsuiji K, Yoshida M, Matsui O. Angiographic evaluation of feeding arteries of hepatocellular carcinoma in the caudate lobe of the liver. *Cardiovasc Intervent Radiol* 2011; 34 (6):1244-1253.
 44. Miyayama S, Yamashiro M, Shibata Y, Hashimoto M, Yoshida M, Tsuiji K, Toshima F, Matsui O. Arterial blood supply to the caudate lobe of the liver from the proximal branches of the right inferior phrenic artery in patients with recurrent hepatocellular carcinoma. *Jpn J Radiol* 2012; 30 (1): 45-52.
 45. Miyayama S, Yamashiro M, Shibata Y, Hashimoto M, Yoshida M, Tsuiji K, Toshima F, Matsui O. Thrombosis and thromboaspiration for acute thromboembolic occlusion in the upper extremity. *Jpn J Radiol* 2012; 30 (2):180-184.
 46. Miyayama S, Yamashiro M, Shibata Y, Hashimoto M, Yoshida M, Tsuiji K, Toshima F, Matsui O. Comparison of local control effects of superselective transcatheter arterial chemoembolization using epirubicin plus mitomycin C and miriplatin for hepatocellular carcinoma. *Jpn J Radiol* 2012; 30 (3):263-270.
 47. Miyayama S, Yamashiro M, Shibata Y, Hashimoto M, Yoshida M, Tsuiji K, Toshima F, Matsui O. Variations in feeding arteries of hepatocellular carcinoma located in the left hepatic lobe. *Jpn J Radiol* 2012; 30 (6):471-479.
 48. Miyayama S, Yamashiro M, Shibata Y, Hashimoto M, Yoshida M, Tsuiji K, Toshima F, Matsui O. Origins of feeding arteries of hepatocellular carcinoma located near the umbilical fissure of the left hepatic lobe: angiographic evaluation. *Cardiovasc Intervent Radiol* 2012; 35 (6):1380-1387.

49. Miyayama S, Yamashiro M, Hashimoto M, Yoshida M, Hashimoto N, Ikuno M, Okumura K, Matsui O. Clinical features of hepatocellular carcinoma supplied by the left internal mammary artery. *Jpn J Radiol* 2012; 30 (10):798-805.
50. Miyayama S, Yamashiro M, Hashimoto M, Hashimoto N, Ikuno M, Okumura K, Yoshida M, Matsui O. Blood supply of the main bile duct from the caudate artery and medial subsegmental artery of the hepatic artery: evaluation using images obtained during transcatheter arterial chemoembolization. *Hepatol Res* 2013;
51. Miyayama S, Yamashiro M, Hashimoto M, Hashimoto N, Ikuno M, Okumura K, Yoshida M, Matsui O. Identification of small hepatocellular carcinoma and tumor-feeding branches with cone-beam CT guidance technology during transcatheter arterial chemoembolization. *J Vasc Interv Radiol* 2013; 24 (4):501-508.
52. Miyayama S, Yamashiro M, Hashimoto M, Hashimoto N, Ikuno M, Okumura K, Matsui O. Hepatojejunostomy obstruction caused by recurrent malignant tumor: percutaneous transhepatic metallic stent with "U-turn" technique. *J Vasc Interv Radiol* 2013; 24 (6):913-915.
53. Miyayama S, Yamashiro M, Hashimoto M, Hashimoto N, Ikuno M, Okumura K, Yoshida M, Matsui O. Comparison of local control in transcatheter arterial chemoembolization of hepatocellular carcinoma ≤ 6 cm with or without intraprocedural monitoring of the embolized area using cone-beam computed tomography. *Cardiovasc Intervent Radiol* 2014; 37 (2):388-395.
54. Miyayama S, Minami T, Kozaka K, Ushioji Y. Small needle puncture of a central venous obstruction in a hemodialysis patient that could not be traversed by a conventional technique. *Cardiovasc Interv Ther* 2014; 29 (3):261-265.
55. Miyayama S, Yamashiro M, Ikuno M, Okumura K, Yoshida M, Kato T, Ushioji Y. Percutaneous transluminal angioplasty of a non-mainstream venous route to restore an occluded hemodialysis fistula. *Jpn J Radiol* 2014; 32 (2):117-122.
56. Guidelines on the use of gelatin sponge particles in embolotherapy. Miyayama S, Yamakado K, Anai H, Abo D, Minami T, Takaki H, Kodama T, Yamanaka T, Nishiofuku H, Morimoto K, Soyama T, Hasegawa Y, Nakamura K, Yamanishi T, Sato M, Nakajima Y. *Jpn J Radiol* 2014; 32 (4):242-250.
57. Miyayama S, Yamashiro M, Ikuno M, Okumura K, Yoshida M. Ultraselective transcatheter arterial chemoembolization for small hepatocellular carcinoma guided by automated tumor-feeders detection software: technical success and short-term tumor response. *Abdom Imaging* 2014; 39 (3):645-656.
58. Miyayama S, Yamashiro M, Ogi T, Kayahashi M, Kawamura K, Yoshida M, Terada

- T, Kosugi I. Usefulness of automated feeder-detection software for identification of access routes to small pancreaticoduodenal artery aneurysms during embolotherapy. *Vascular* 2015; 23 (6):663-667.
59. Miyayama S, Terada T, Tamaki M. Ruptured pancreaticoduodenal artery aneurysm associated with median arcuate ligament compression and aortic dissection successfully treated with embolotherapy. *Ann Vasc Dis* 2015; 8 (1):40-42.
 60. Miyayama S, Yamashiro M, Nagai K, Tohyama J, Kawamura K, Yoshida M, Sakuragawa N. Efficacy of automated tumor-feeder detection software using cone-beam computed tomography technology in transarterial embolization through extrahepatic collateral vessels for malignant hepatic tumors. *Hepatol Res* 2016; 46 (2):166-173.
 61. Miyayama S, Yamashiro M, Nagai K, Tohyama J, Kawamura K, Yoshida M, Sakuragawa N. Evaluation of tumor recurrence after superselective conventional transcatheter arterial chemoembolization for hepatocellular carcinoma: comparison of computed tomography and gadoxetate disodium-enhanced magnetic resonance imaging. *Hepatol Res* 2016; 46 (9):890-898.
 62. Miyayama S, Yamashiro M, Nagai K, Tohyama J, Kawamura K. Efficacy of automated tumor-feeder detection software using cone-beam computed tomography technology in transarterial targeted therapy. *Interventional Radiology* 2016; 1 (1):28-38.
 63. Miyayama S, Matsui O. Superselective conventional transarterial chemoembolization for hepatocellular carcinoma: rationale, technique, and outcome. *J Vasc Interv Radiol* 2016; 27 (9):1269-1278.
 64. Miyayama S, Yamashiro M, Nagai K, Yokka A, Yoshida M, Sakuragawa N. Performance of novel virtual parenchymal perfusion software visualizing embolized areas of transcatheter arterial chemoembolization for hepatocellular carcinoma. *Hepatol Res*. 2017; 47 (5):446-454.
 65. Miyayama S, Yamashiro M, Nagai K, Yokka A. Use of a catheter with a large side hole or cleft in selective catheterization of small branch arteries. *Interventional Radiology* 2016; 1 (2):67-74.
 66. Miyayama S, Yamashiro M, Yagai K, Yokka A, Toya D. Long-term patency of metallic stents in a patient with bile duct injury refractory to long-term catheter placement following right hepatectomy for liver metastasis. *Interventional Radiology* 2017; 2 (2):10-13.
 67. Miyayama S, Yamashiro M, Nagai K, Yokka A, Yoshida M, Sakuragawa N. Excretion of necrotic hepatocellular carcinoma tissues into the biliary system after

- transcatheter arterial chemoembolization. *Hepatol Res* 2017; 47 (13):1390-1396.
68. Miyayama S, Yamashiro M, Nagai K, Sugimori N, Iwata K, Sakuragawa N. Usefulness of dual-phase cone-beam computed tomography during arteriography and automated tumour-feeder detection software in transarterial embolization for obscure arterial bleeding in the abdomen. *J Med Imaging Radiat Oncol* 2018; 62 (3):364-369.
69. Miyayama S, Yamashiro M, Sugimori N, Ikeda R, Okimura K, Sakuragawa N. Outcomes of patients with hepatocellular carcinoma treated with conventional transarterial chemoembolization using guidance software. *J Vasc Interv Radiol* 2018 (in press).

Book (in English)

1. Miyayama S. Transcatheter arterial embolization. In: Ohkochi N ed. *Therapy for hepatocellular carcinoma – Etiology and treatment –*. New York, USA: Nova Biomedical, 2014; 159-243.